

Efficient ion acceleration by collective laser-driven electron dynamics with ultra-thin foil targets

S. STEINKE,¹ A. HENIG,^{2,3} M. SCHNÜRER,¹ T. SOKOLLIK,¹ P.V. NICKLES,^{1,4} D. JUNG,^{3,5}
D. KIEFER,^{2,3} R. HÖRLEIN,^{2,3} J. SCHREIBER,^{2,3,6} T. TAJIMA,^{3,7} X.Q. YAN,^{2,8} M. HEGELICH,^{3,5}
J. MEYER-TER-VEHN,² W. SANDNER,¹ AND D. HABS^{3,2}

¹Max-Born-Institut, D-12489 Berlin, Germany

²Max-Planck-Institut f. Quantenoptik, D-85748 Garching, Germany

³Fakultät f. Physik, LMU München, D-85748 Garching, Germany

⁴Gwangju Institute of Science and Technology, GIST, Gwangju 500-712, Republic of Korea

⁵Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA

⁶Imperial College London, SW7 2BZ, UK

⁷Photomedical Research Center, JAEA, Kyoto, Japan

⁸State Key Lab of Nuclear physics and technology, Peking University, 100871, Beijing, China

(RECEIVED 24 November 2009; ACCEPTED 29 January 2010)

Abstract

Experiments on ion acceleration by irradiation of ultra-thin diamond-like carbon (DLC) foils, with thicknesses well below the skin depth, irradiated with laser pulses of ultra-high contrast and linear polarization, are presented. A maximum energy of 13 MeV for protons and 71 MeV for carbon ions is observed with a conversion efficiency of $\sim 10\%$. Two-dimensional particle-in-cell (PIC) simulations reveal that the increase in ion energies can be attributed to a dominantly collective rather than thermal motion of the foil electrons, when the target becomes transparent for the incident laser pulse.

Keywords: Laser-driven acceleration; Laser-plasma interactions; Laser-produced plasma; Particle-in-cell method; Plasma simulation

INTRODUCTION

Recent experiments in the field of relativistic laser-plasma interaction have shown that the conversion efficiency (CE) from the laser to the kinetic energy of the ion bunch as well as the maximal energy of the ions can be improved by the use of ultra-thin foil targets in the range of 100 nm (Andreev *et al.*, 2009; Antici *et al.*, 2007; Ceccotti *et al.*, 2007; Limpouch *et al.*, 2008; Neely *et al.*, 2006) or by a special target geometry (Schnuerer *et al.*, 2005; Wang *et al.*, 2009). Also, spectral shaping of the ion bunch was demonstrated (Flippo *et al.*, 2007; Schwoerer *et al.*, 2006; Ter-Avetisyan *et al.*, 2006, 2008).

Additionally, simulations emphasized that besides the standard target normal sheath acceleration (TNSA) (Hatchett *et al.*, 2000; Wilks *et al.*, 1992) other, radiation pressure dominated acceleration mechanisms become possible for ultra-thin targets (Klimo *et al.*, 2008; Robinson *et al.*, 2008; Yan *et al.*, 2008). An ultra-high contrast is required to avoid substantial

expansion of the targets before the interaction with the main pulse. These conditions can be achieved by the use of a specially designed laser system in combination with double plasma mirrors (DPM) (Andreev *et al.*, 2009; Levy *et al.*, 2007; Wittmann *et al.*, 2006). The partial transmission of the intense laser pulse through an expanding target plays a decisive role (Albright *et al.*, 2007; d’Humieres *et al.*, 2005; Yin *et al.*, 2006) and has been lately experimentally demonstrated with circularly polarized laser pulses (Henig *et al.*, 2009b).

In this article, we present experimental results on ion acceleration from DLC foils of thicknesses ranging from 50 nm down to 2.9 nm. The targets are irradiated by linear polarized pulses of 45 fs (FWHM) duration focused to a peak intensity of up to 5×10^{19} W/cm². We find an optimum in ion acceleration at a target thickness of 5.6 nm, where ion energies reach values of 13 MeV for protons and 71 MeV for carbon ions. Thus, for the ultra-short pulses used, the optimum foil thickness is well below the collisional skin depth $l_s \sim 13$ nm of the heated target, which is thus becoming transparent to the laser. The corresponding CE is reaching values of $\sim 1.6\%$ for protons ($E > 2$ MeV) and $\sim 10\%$ in the case of

Address correspondence and reprint requests to: S. Steinke, Max-Born-Institut, D-12489 Berlin, Germany. E-mail: steinke@mbi-berlin.de

C^{6+} ($E > 5$ MeV). By means of 2D PIC-simulations, we find that the transmitted laser field imposes a dominantly collective rather than thermal motion of the foil electrons, leading to increased ion energies and enhanced CE. Our findings are supported by a semi-analytical model, which shows good agreement with the experimental results.

EXPERIMENT

The experiments were performed at the MBI-TW Ti:sapph laser of central wavelength 810 nm delivering 1.2 J in 45 fs FWHM pulses with an amplified spontaneous emission (ASE) contrast ratio smaller than 10^{-7} up to ~ 10 ps prior to the arrival of the main peak. By means of a re-collimating DPM (Andreev *et al.*, 2009), this contrast was increased by estimated four orders of magnitude (Levy *et al.*, 2007), which is essential for the suppression of pre-heating and expansion due to the pulse background. The energy throughput this DPM system was improved to values of 60–65% compared to our previous experiments (Andreev *et al.*, 2009), resulting in pulse energies of 0.7 J. Finally, the laser pulse was focused on the DLC target with a $f/2.5$ parabolic mirror down to $6 \mu\text{m}$ diameter and is diffraction limited by 30% under normal incidence. This corresponds to a peak intensity of $2.6 \times 10^{19} \text{ W/cm}^2$ or a normalized vector potential of $a_L = 3.6$. For a second set of experiments, the focusing procedure was improved, leading to a focus diameter of $3.6 \mu\text{m}$ and therefore a peak intensity of $5 \times 10^{19} \text{ W/cm}^2$ or $a_L = 5$. The resulting ions were detected with a MCP coupled to a Thomson-Parabola (Ter-Avetisyan *et al.*, 2005). Additionally, the transmitted laser pulse was registered with a 12-bit optical grating spectrometer.

DLC targets of thicknesses ranging from 2.9–50 nm were used, having a density of 2.7 g/cm^3 . Owing to the high fraction of sp^3 -, i.e., diamond-like bonds of $\sim 75\%$, DLC offers unique properties for the production of mechanically stable, ultra-thin, free standing targets, such as exceptionally high tensile strength, hardness and heat resistance. The thickness of the DLC foils was characterized by means of an atomic force microscope (AFM), including the hydrocarbon contamination layer on the target surface, which was present during the experiments. In addition, in order to precisely determine the structure of the contamination layer, the depth-dependent composition of the foil was measured via elastic recoil detection analysis (ERDA). From these measurements we obtain a thickness of ~ 1 nm for the hydrocarbon contamination layer. Throughout the manuscript we are referring to the combined thickness of bulk and surface layer as it appears in the actual ion acceleration experiment presented.

The maximum detectable energy/nucleon (E_{max}) obtained for protons and fully ionized carbon ions C^{6+} is plotted as a function of the target thickness in Figure 1 for two different laser intensities. The cut-off energies for both ion species exhibit a strong dependence on target thickness. In case of protons and $a_0 = 5$, E_{max} increases from around 7.5 MeV for a 40 nm foil up to 13 MeV for a 5.6 nm foil, while for




Fig. 1. (Color online) Maximum energy per atomic mass unit for both protons and carbon ions plotted vs. foil thickness for values of the normalized laser vector potential of $a_0 = 3.6$ and $a_0 = 5$. The experimental results are in excellent agreement with numbers deduced from 2D-PIC-simulations.

C^{6+} the maximum energy rises from 26 MeV for use of a 40 nm foils to 71 MeV for 5.6 nm. Further decrease of the target thickness down to 2.9 nm results in a steep drop of the observed ion energies. The energy distributions of all species are continuous as expected. However, especially the observed carbon C^{6+} energy of 71 MeV reaches for the first time a range of values that were previously only accessible by large single shot Nd: glass laser systems with 30–50 J pulse energy (Henig *et al.*, 2009a; Krushelnick *et al.*, 2005). The shot-to-shot variations of 10%, indicated by the error bars, arise mainly from fluctuations of the laser pulse itself and from macroscopic modulations of the target surface.




Fig. 2. (Color online) Calculated CE for protons and C^{6+} ions as a function of the target thickness, based on an energy dependent divergence angle of the ion beam, which was extracted from PIC simulations and supported by measured beam profiles (cf. figure inset). The experimental data are in good agreement with the values of the CE obtained from PIC simulations.

The CE (Fig. 2) was calculated by numerically convoluting an energy dependent divergence of the accelerated ion beam with the initially measured ($a_0 = 5$) spectra. These values (cf. inset in Fig. 2) were extracted from PIC simulations and supported by experimentally obtained proton beam profiles using a stack of radiochromic film (RCF) layers. Comparing those to the divergencies of ion beams generated by other laser systems (Nurnberg *et al.*, 2009), our divergence is smaller and the linear dependence on the energy is reasonable in the considered high-energy part. The dependence of the CE on the target thickness is showing analogous characteristics to E_{\max} (cf. Fig. 1). A sudden rise is observed when entering the laser transparency regime ($D < l_s$) reaching 10.5% for C^{6+} and 1.6% for protons at the optimum thickness. The CE drops down to below 1% when the target thickness reaches the TNSA dominated regime (Fuchs *et al.*, 2006; Neely *et al.*, 2006).

Furthermore, the transmitted laser pulse was imaged to a 12-bit optical grating spectrometer for several target thicknesses. Numerical integrations of the measured spectra, referenced to a shot without target yield to the transmittance T of the DLC targets. In Figure 3, these values for the transmittance are plotted as a function of the target thickness and compared to the analytical formula

$$T \cong 1/[1 + (\pi\sigma/\gamma_p)^2], \quad (1)$$

where γ_p is the Lorentz factor in the laser $\sqrt{a_L^2 + 1}$ (Vshivkov *et al.*, 1998). The experimental results are in agreement with the analytical model.

SIMULATION

The experiments were compared to 2D-PIC simulations, where the laser pulse was modeled by a Gaussian shape in




Fig. 3. Measured laser pulse ($a_0 = 5$) transmittance of the DLC targets as a function of the target thickness compared to the analytical model of Vshivkov *et al.* (1998), Eq. (1).

time with a FWHM of 16 laser cycles, a Gaussian intensity distribution in focus with a FWHM spot size of $4 \mu\text{m}$ and an $a_0 = 5$ ($a_0 = 3.6$). It interacts with a rectangular shaped plasma of initial density $n_e = 500 \times n_c$ consisting of 90% carbon ions and 10% protons (in number density) to account for the presence of a contamination layer. The simulation box is composed of 1200×10000 cells with 1000 particles per cell and a total size of $(10 \times 20) \mu\text{m}^2$. The total simulation time τ given in laser cycles is 120.

Figure 1 shows that the simulated maximum proton energies as well as the carbon energies are in excellent agreement with the experimental data. In particular, the optimum target thickness of 5.6 nm is reproduced. This thickness for the peak ion energy is consistent with the thickness given by the empirical relation $\sigma \approx 3 + 0.4 \times a_0 = 5$ that was found in multiparametric PIC-simulation studies by Esirkepov *et al.* (2006). Here, a normalized areal electron density

$$\sigma = (n_e/n_c) \cdot (D/\lambda_L), \quad (2)$$

is introduced, where D is the target thickness, λ_L is the laser wave length, n_e is the electron density of the target, and n_c is the critical electron density, which in our case calculates to $\sigma = 4.6$. For foil thicknesses below the optimum ($\sigma < 3 + 0.4 \times a_0$), the plasma becomes increasingly transparent (Fig. 3) and the pulse is more transmitted than absorbed. Due to the low number of electrons in the focal volume ($\sim 10^{11}$) their electric current is no longer sufficient to (1) reflect the laser pulse and (2) to establish an effective longitudinal charge separation field. This results in a sudden drop in ion energies and 50% of the CE (the same amount as the reduction of the target thickness), as it was observed in the experiment (see Figs. 1, 2). In case of $\sigma > 3 + 0.4 \times a_0$, the laser intensity is not sufficient to generate the maximum possible displacement of all electrons within the focal volume which gives rise to a decrease in the longitudinal charge separation field. Note that the optimum thickness for ion acceleration predicted theoretically and observed experimentally is much smaller than previously used target thicknesses.

In Figure 4a, the observed carbon energy spectrum is shown along with the simulated one agreeing with each other in their overall spectral shape. To illustrate the underlying ion acceleration mechanism in our case, we show in Figures 4c and 4d detailed electron and carbon densities after 40 laser cycles as a function of the laser propagation direction x and the transverse coordinate y in units of the laser wave length λ_L . It is immediately striking that ions are accelerated strongly asymmetric heavily favoring the direction of laser propagation. This is in strong contrast to the model based on the self-consistent solution of the Poisson equation presented in Andreev *et al.* (2008) which predicts a symmetric acceleration that happens primarily after the end of the laser pulse. From Figures 4c and 4d it can be seen that the carbon ions are accompanied by co-moving electrons, which accelerate the ions in forward direction.




Fig. 4. (Color online) (a) Comparison between experimentally obtained and simulated C^{6+} spectrum as deduced from a 2D-PIC simulation of a 4.5 nm foil target (b) Electron propagation angle (blue circles) and laser transverse electric field E_y (red line), plotted vs. laser propagation direction x , given in units of the laser wavelength λ . τ is given in laser cycles. The green line at $x = 5\lambda$ gives the initial location of the target. For the same simulation, (c) gives the electron and (d) the carbon density distribution vs. x at time $t = 40\tau$.

Unlike the TNSA consideration in case of thicker targets, in our current case, coherence of laser-driven electron motion manifests importantly. In the typical TNSA with thicker targets, none of the laser field is transmitted as the foil remains opaque throughout the interaction duration. The electron motion is an indirect result of interaction with the laser at the front surface of the target and subsequent multiple interactions with matter in the target bulk. That scenario leads to thermal motion as the electrons traverse the target to set up a quasi-static field at the back. In contrast, in nm-scale foils of thickness below the skin depth the laser imprints a directed motion onto electrons as those targets are partially transparent. These coherent electron motions can not be treated as thermal motion, but rather as collective motion tied to the laser and thus directly to the laser amplitude a_0 . The propagation angle of electrons upon laser interaction with the target as extracted from the 2D PIC simulation at time 30τ is shown in Figure 4b. Obviously, the electron dynamics are classified in three categories. (1) electrons are driven forward by laser, (2) electrons are forced to return backwards due to the electrostatic field, (3) electrons are performing a collective, transversal oscillation owing to the laser-created

coherent wave structure. Additionally, the laser transmission coefficient can be determined to be $T \sim 0.1$ which agrees with the theoretical given by Eq. (1).

ANALYTICAL MODEL

We compared our results to a number of analytical scaling laws that have been derived for the standard scenario of ion acceleration from the back surface of a thin, but opaque foil, i.e., target normal sheath acceleration (TNSA) (Andreev *et al.*, 2008; Fuchs *et al.*, 2006; Schreiber *et al.*, 2006). However, we find that all of the existing analytical models fail to even qualitatively predict the two main features seen experimentally, namely the distinct peak in ion energies at $\sigma \approx 3 + 0.4 \times a_0$ as well as the strong dependence of the cut-off value on target thickness. Consequently, a refined theory is needed.

Initially, without assuming thermalized electrons (i.e., in contrast to Mora (2003)), we now formulate the maximal ion energies obtained in the laser driven foil interaction in a semi-analytical model based on Mako and Tajima (1984).

Following the analysis of Yan *et al.* (2009), the forward current density of electrons J and electron density n_e are related through

$$J = -e \int_v^{V_{\max}} V_x g dV_x, \quad (3)$$

and

$$n_e = \frac{2}{e} \int_0^{V_{\max}} \frac{dJ/dv}{v} dv, \quad (4)$$

where g is the electron distribution function and $V_{\max} = c\sqrt{1 - m_e^2 c^4 / (E_0 + e\phi)^2}$. Here we note that at a given position the electron energy E_0 in the electrostatic potential ϕ is given as an implicit function of space and time through $\varepsilon = \gamma m_e c^2 - e\phi$. We find that

$$J(\varepsilon) = -J_0(1 - \varepsilon/E_0)^\alpha, \quad (5)$$

where α , a measure for the coherence is found to be about 3 in our simulation for a target thickness of 4.5 nm. Considering the relativistic dynamics of electrons in our regime, the electron density as a function of the current density may be integrated to yield

$$\begin{aligned} n_e &= \frac{2}{e} \int_0^{V_{\max}} \frac{dJ/dv}{v} dv = \frac{2}{ec} \int_{-e\phi}^{E_{\max}} \frac{dJ}{d\varepsilon} d\varepsilon \\ &= \frac{2J_0}{ec} \left(1 + \frac{e\phi}{E_0}\right)^\alpha. \end{aligned} \quad (6)$$

The integration of the nonlinear coupled equations for nonrelativistic ions under the self-consistently evolving potential using Eq. (6) with the boundary condition and initial value settings of Mako and Tajima (1984) leads to the maximum ion (with charge Q) energy as $\varepsilon_{\max} = (2\alpha + 1)QE_0$, i.e. the maximum carbon energy is determined by the maximum electron kinetic energy. In the present experimental condition we recognize that the electron kinetic energy is primarily dominated by the laser driven collective momentum so that

$$\varepsilon_{\max} = (2\alpha + 1)Qmc^2(\gamma_p - 1), \quad (7)$$

which gives rise to $(2\alpha + 1)Qmc^2(\sqrt{a_L^2 + 1} - 1)$. When we insert the exponent of $\alpha = 3$ as observed in our 2D-PIC simulations, we obtain for the case of $a_0 = 3.6$ a theoretical maximum carbon (proton) energy of 59 MeV (9.6 MeV), in the case of $a_0 = 5$, 88 MeV (14.3 MeV). This is in agreement with the experimental value between 36-71 MeV (11–13 MeV).

Again, for conventional TNSA we do not expect a strong dependence of the maximum ion energy on target thickness when the target is much smaller than the transversal size of the laser focal spot as it is the case in the present study. The thickness dependence in TNSA is mainly due to

transversal spread of the laser accelerated electrons on their way through the target (Nickles *et al.*, 2007; Schreiber *et al.*, 2006).

Instead, the strong increase in ε_{\max} as it was observed in the experiment is related to the growing transmission for thinner targets leading to collective electron dynamics imposed by the laser field. For thick targets, however, α drops towards zero and Eq. (7) reveals the dependence of ε_{\max} on the hot electron energy $mc^2(\sqrt{a_L^2 + 1} - 1)$ (Mora, 2003; Wilks *et al.*, 1992).

SUMMARY

In summary, ion acceleration from ultra-thin DLC foils of thickness 2.9–50 nm was studied. A strong dependence of the resulting maximum energies on target thickness was observed experimentally, with a pronounced optimum for an initial foil extension of 5.6 nm with a very high conversion efficiency from the laser to the kinetic energy of the ion bunch. Here, proton energies of 13 MeV with a CE of 1.6% and carbon ions C^{6+} of 71 MeV with a CE of 10% were obtained with laser pulse energies as low as 0.7 J. Our experimental results are in excellent agreement with 2D-PIC simulations. However, we find that previously published TNSA scaling laws based on analytical models (Andreev *et al.*, 2008; Fuchs *et al.*, 2006; Schreiber *et al.*, 2006) fail to interpret our results. Therefore, a new semi-analytical model based on Mako and Tajima (1984) was developed, showing good agreement with the experimental data.

Demonstrating enhanced ion energies and CE from tabletop laser systems operating at high repetition rates represents a major step towards potential applications such as medical or industrial motivated approaches.

ACKNOWLEDGMENTS

This work was partly supported by Deutsche Forschungsgemeinschaft through Transregio SFB TR18 and the DFG-Cluster of Excellence Munich-Centre for Advanced Photonics (MAP). A. Henig, D. Kiefer and D. Jung acknowledge financial support from IMPRS-APS, J. Schreiber from DAAD, X. Q. Yan from the Humboldt foundation and NSFC(10855001).

REFERENCES

- ALBRIGHT, B.J., YIN, L., BOWERS, KEVIN J., HEGELICH, B.M., FLIPPO, K.A., KWAN, T.J.T. & FERNANDEZ, J.C. (2007). Relativistic buneman instability in the laser breakout afterburner. *Phys. Plasmas* **14**, 094502–094504.
- ANDREEV, A., LEVY, A., CECCOTTI, T., THAURY, C., PLATONOV, K., LOCH, R.A. & MARTIN, P.H. (2008). Fast-ion energy-flux enhancement from ultrathin foils irradiated by intense and high-contrast short laser pulses. *Phys. Rev. Lett.* **101**, 155002.
- ANDREEV, A.A., STEINKE, S., SOKOLLIK, T., SCHNUEERER, M., TER AVETSIYAN, S., PLATONOV, K.Y. & NICKLES, P.V. (2009). Optimal ion acceleration from ultrathin foils irradiated by a

- profiled laser pulse of relativistic intensity. *Phys. Plasmas* **16**, 013103.
- ANTICI, P., FUCHS, J., D'HUMIERES, E., LEFEBVRE, E., BORGHESI, M., BRAMBRINK, E., CECCHETTI, C.A., GAILLARD, S., ROMAGNANI, L., SENTOKU, Y., TONCIAN, T., WILLI, O., AUDEBERT, P. & PEPIN, H. (2007). Energetic protons generated by ultra-high contrast laser pulses interacting with ultrathin targets. *Phys. Plasmas* **14**, 030701.
- CECCOTTI, T., LEVY, A., POPESCU, H., REAU, F., D'OLIVEIRA, P., MONOT, P., GEINDRE, J.P., LEFEBVRE, E. & MARTIN, P. (2007). Proton acceleration with high-intensity ultrahigh-contrast laser pulses. *Phys. Rev. Lett.* **99**, 185002.
- D'HUMIERES, Emmanuel, LEFEBVRE, Erik, GREMILLET, Laurent & MALKA, Victor. (2005). Proton acceleration mechanisms in high-intensity laser interaction with thin foils. *Phys. Plasmas* **12**, 062704–062713.
- ESIRKEPOV, T., YAMAGIWA, M. & TAJIMA, T. (2006). Laser ion-acceleration scaling laws seen in multiparametric particle-in-cell simulations. *Phys. Rev. Lett.* **96**, 105001.
- FLIPPO, K., HEGELICH, B.M., ALBRIGHT, B.J., YIN, L., GAUTIER, D.C., LETZRING, S., SCHOLLMEIER, M., SCHREIBER, J., SCHULZE, R., FERN, Aacute & NDEZ, J.C. (2007). Laser-driven ion accelerators: Spectral control, monoenergetic ions and new acceleration mechanisms. *Laser Part. Beams* **25**, 3–8.
- FUCHS, J., ANTICI, P., D'HUMIERES, E., LEFEBVRE, E., BORGHESI, M., BRAMBRINK, E., CECCHETTI, C.A., KALUZA, M., MALKA, V., MANCLOSSI, M., MEYRONEINC, S., MORA, P., SCHREIBER, J., TONCIAN, T., PEPIN, H. & AUDEBERT, R. (2006). Laser-driven proton scaling laws and new paths towards energy increase. *Nat. Phys.* **2**, 48–54.
- HATCHETT, S.P., BROWN, C.G., COWAN, T.E., HENRY, E.A., JOHNSON, J.S., KEY, M.H., KOCH, J.A., LANGDON, A.B., LASINSKI, B.F., LEE, R.W., MACKINNON, A.J., PENNINGTON, D.M., PERRY, M.D., PHILLIPS, T.W., ROTH, M., SANGSTER, T.C., SINGH, M.S., SNAVELY, R.A., STOYER, M.A., WILKS, S.C. & YASUIKE, K. (2000). Electron, photon, and ion beams from the relativistic interaction of petawatt laser pulses with solid targets. *Phys. Plasmas* **7**, 2076–2082.
- HENIG, A., KIEFER, D., MARKEY, K., GAUTIER, D.C., FLIPPO, K.A., LETZRING, S., JOHNSON, R.P., SHIMADA, T., YIN, L., ALBRIGHT, B.J., BOWERS, K.J., FERNANDEZ, J.C., RYKOVANOV, S.G., WU, H.C., ZEPF, M., JUNG, D., LIECHTENSTEIN, V.K., SCHREIBER, J., HABS, D. & HEGELICH, B.M. (2009a). Enhanced laser-driven ion acceleration in the relativistic transparency regime. *Phys. Rev. Lett.* **103**, 045002.
- HENIG, A., STEINKE, S., SCHNUEERER, M., SOKOLLIK, T., HORLEIN, R., KIEFER, D., JUNG, D., SCHREIBER, J., HEGELICH, B.M., YAN, X.Q., MEYER-TER VEHN, J., TAJIMA, T., NICKLES, P.V., SANDNER, W. & HABS, D. (2009b). Radiation-pressure acceleration of ion beams driven by circularly polarized laser pulses. *Phys. Rev. Lett.* **103**, 245003.
- KLIMO, O., PSIKAL, J., LIMPOUCH, J. & TIKHONCHUK, V.T. (2008). Monoenergetic ion beams from ultrathin foils irradiated by ultrahigh-contrast circularly polarized laser pulses. *Phys. Rev. Lett.* **111**, 031301.
- KRUSHELNICK, K., CLARK, E.L., BEG, F.N., DANGOR, A.E., NAJMUDIN, Z., NORREYS, P.A., WEI, M. & ZEPF, M. (2005). High intensity laser-plasma sources of ions-physics and future applications. *Plasma Phys. Contr. Fusion* **47**, B451–B463.
- LEVY, A., CECCOTTI, T., D'OLIVEIRA, P., REAU, F., PERDRIX, M., QUERE, F., MONOT, P., BOUGEARD, M., LAGADEC, H., MARTIN, P., GEINDRE, J.P. & AUDEBERT, P. (2007). Double plasma mirror for ultrahigh temporal contrast ultraintense laser pulses. *Opt. Lett.* **32**, 310–312.
- LIMPOUCH, J., PSIKAL, J., ANDREEV, A.A., PLATONOV, K. Yu & KAWATA, S. (2008). Enhanced laser ion acceleration from mass-limited targets. *Laser Part. Beams* **26**, 225–234.
- MAKO, F. & TAJIMA, T. (1984). Collective ion-acceleration by a reflexing electron-beam – model and scaling. *Phys. Fluids* **27**, 1815–1820.
- MORA, P. (2003). Plasma expansion into a vacuum. *Phys. Rev. Lett.* **90**, 185002.
- NEELY, D., FOSTER, P., ROBINSON, A., LINDAU, F., LUNDH, O., PERSSON, A., WAHLSTROM, C.G. & MCKENNA, P. (2006). Enhanced proton beams from ultrathin targets driven by high contrast laser pulses. *Appl. Phys. Lett.* **89**, 021502.
- NICKLES, P.V., TER-AVETISYAN, S., SCHNUEERER, M., SOKOLLIK, T., SANDNER, W., SCHREIBER, J., HILSCHER, D., JAHNKE, U., ANDREEV, A. & TIKHONCHUK, V. (2007). Review of ultrafast ion acceleration experiments in laser plasma at max born institute. *Laser Part. Beams* **25**, 347–363.
- NURNBERG, F., SCHOLLMEIER, M., BRAMBRINK, E., BLAZEVIC, A., CARROLL, D.C., FLIPPO, K., GAUTIER, D.C., GEISSEL, M., HARRES, K., HEGELICH, B.M., LUNDH, O., MARKEY, K., MCKENNA, P., NEELY, D., SCHREIBER, J. & ROTH, M. (2009). Radiochromic film imaging spectroscopy of laser-accelerated proton beams. *Rev. Sci. Instru.* **80**, 033301–033313.
- ROBINSON, A.P.L., ZEPF, M., KAR, S., EVANS, R.G. & BELLEI, C. (2008). Radiation pressure acceleration of thin foils with circularly polarized laser pulses. *New J. Phys.* **10**, 013021.
- SCHNUEERER, M., TER-AVETISYAN, S., BUSCH, S., RISSE, E., KALACHNIKOV, M.P., SANDNER, W. & NICKLES, P.V. (2005). Ion acceleration with ultrafast laser driven water droplets. *Laser Part. Beams* **23**, 337–343.
- SCHREIBER, J., BELL, F., GRUENER, F., SCHRAMM, U., GEISLER, M., SCHNUEERER, M., TER-AVETISYAN, S., HEGELICH, B.M., COBBLE, J., BRAMBRINK, E., FUCHS, J., AUDEBERT, P. & HABS, D. (2006). Analytical model for ion acceleration by high-intensity laser pulses. *Phys. Rev. Lett.* **97**, 045005.
- SCHWOERER, H., PFOTENHAUER, S., JACKEL, O., AMTHOR, K.U., LIESFELD, B., ZIEGLER, W., SAUERBREY, R., LEDINGHAM, K.W.D. & ESIRKEPOV, T. (2006). Laser-plasma acceleration of quasi-monoenergetic protons from microstructured targets. *Nat.* **439**, 445–448.
- TER-AVETISYAN, S., SCHNUEERER, M., NICKLES, P.V., KALASHNIKOV, M., RISSE, E., SOKOLLIK, T., SANDNER, W., ANDREEV, A. & TIKHONCHUK, V. (2006). Quasimonoenergetic deuteron bursts produced by ultraintense laser pulses. *Phys. Rev. Lett.* **96**, 145006.
- TER-AVETISYAN, S., SCHNUEERER, M., POLSTER, R., NICKLES, P.V. & SANDNER, W. (2008). First demonstration of collimation and monochromatisation of a laser accelerated proton burst. *Laser Part. Beams* **26**, 637–642.
- TER-AVETISYAN, S., SCHNUEERER, M. & NICKLES, P.V. (2005). Time resolved corpuscular diagnostics of plasmas produced with high-intensity femtosecond laser pulses. *J. Phys. D-Appl. Phys.* **38**, 863–867.
- VSHIVKOV, V.A., NAUMOVA, N.M., PEGORARO, F. & BULANOV, S.V. (1998). Nonlinear electrodynamics of the interaction of ultraintense laser pulses with a thin foil. *Phys. Plasmas* **5**, 2727–2741.

- WANG, X., YU, W., YU, M.Y., SENECHA, V.K., XU, H., WANG, J.W., YUAN, X. & SHENG, Z.M. (2009). Efficient acceleration of a small dense plasma pellet by consecutive action of multiple short intense laser pulses. *Laser Part. Beams* **27**, 629–634.
- WILKS, S.C., KRUEER, W.L., TABAK, M. & LANGDON, A.B. (1992). Absorption of ultra-intense laser pulses. *Phys. Rev. Lett.* **69**, 1383.
- WITTMANN, T., GEINDRE, J.P., AUDEBERT, P., MARJORIBANKS, R.S., ROUSSEAU, J.P., BURG, F., DOUILLET, D., LEFROU, T., PHUOC, K.T. & CHAMBARET, J.P. (2006). Towards ultrahigh-contrast ultra-intense laser pulses-complete characterization of a double plasma-mirror pulse cleaner. *Rev. Sci. Instru.* **77**, 083109.
- YAN, X.Q., LIN, C., SHENG, Z.M., GUO, Z.Y., LIU, B.C., LU, Y.R., FANG, J.X. & CHEN, J.E. (2008). Generating high-current mono-energetic proton beams by a circularly polarized laser pulse in the phase-stable acceleration regime. *Phys. Rev. Lett.* **100**, 135003.
- YAN, X.Q., TAJIMA, T., HEGELICH, M., YIN, L. & HABS, D. (2009). Theory of laser ion acceleration from a foil target of nanometer thickness. *Appl. Phys. B*, DOI 0.1007/s00340-00009-03707-00345.
- YIN, L., ALBRIGHT, B.J., HEGELICH, B.M. & FERNANDEZ, J.C. (2006). GeV laser ion acceleration from ultrathin targets: The laser break-out afterburner. *Laser Part. Beams* **24**, 291–298.